

# Cotton Variety Guide | 2017



**UT Cotton Agronomy**  
Department of Plant Sciences  
University of Tennessee



Tyson B. Raper, Cotton and Small Grains Specialist  
Department of Plant Sciences

## Contributing Authors

Ryan H. Blair	Extension Area Specialist	UT Extension
Shawn Butler	Research Associate	Department of Plant Sciences
Dalton McCurley	Research Specialist	Department of Plant Sciences
Randi C. Dunagan	Research Associate	West TN Research and Education Center
J. Richard Buntin	Ext Agent III & Director, Crockett Co	UT Extension
Philip W. Shelby	Ext Agent III, Gibson Co	UT Extension
Jake Mallard	Extension Agent II, Madison Co	UT Extension

## 2016 Tennessee Cotton Variety Trials

The University of Tennessee Cotton Agronomy Program provides an unbiased evaluation of experimental and commercial varieties available for production in Tennessee each year. The 2016 program consisted of three types of trials: the Official Variety Trials (OVTs), large replicated on-farm variety trials, and the County Standard Trials (CSTs). The OVTs are small plot, replicated variety trials composed of experimental and commercial varieties. The large replicated on-farm trials and CSTs are large plot variety trials located throughout the Western and Central regions of Tennessee and are only composed of major commercial cultivars. Six OVTs, four large replicated trials, and 15 CSTs were conducted during the 2016 season (Fig.1).



**Figure 1:** County map of Tennessee, with counties participating in the 2016 Large-Plot Variety Trial Program highlighted in orange. Red points represent planted locations of the 2016 Official Variety Trials.

Samples from each plot were ginned at the University of Tennessee Cotton MicroGin located at the West Tennessee AgResearch and Education Center in Jackson, Tennessee. This is a 20-saw gin equipped with a stick machine, inline cleaners and two lint cleaners. No heat was applied at ginning. Lint yields on a per plot basis were calculated from gin turnouts and harvested plot areas. A subsample of lint from each ginned sample was submitted to the USDA Cotton Classing Office in Memphis, Tennessee, for HVI analysis.

Information reported in this publication includes average gin turnout, lint yield, and fiber quality averaged by program. This brief report serves as a precursor for the 2016 Tennessee Cotton Variety Trial Results (PB 1742). Specific results from each trial location and plant growth measurements are included in PB 1742.

This publication is intended to help cotton producers identify varieties that are high yielding, stable in yield performance across environments and years, and consistently produce high quality fiber; therein, included information should also provide those in the seed industry, crop consultants, and UT Extension insight into varietal adaptation of all tested varieties to Tennessee field environments.

# Large On-Farm Trial Results

## County Standard Trials

Lint yield, gin turnout, fiber quality, and CCC loan value of 15 entries entered in the 2016 Tennessee County Standard Trial Program.\*

Yield Rank	Variety	Lint Yield (lb/ac)	Turnout (%)	Mic	Length (in.)	Strength (g/tex)	Unif. (%)	HVI Color	Leaf Grade	Loan Value (¢/lb)
1	PHY 312 WRF	1325 <sub>a</sub>	38.4 <sub>efgh</sub>	4.7 <sub>fgh</sub>	1.20 <sub>bc</sub>	32.0 <sub>cd</sub>	82.7 <sub>abc</sub>	41	4 <sub>abc</sub>	54.75
2	DP 1614 B2XF	1281 <sub>ab</sub>	40.1 <sub>ab</sub>	5.2 <sub>a</sub>	1.20 <sub>bc</sub>	31.4 <sub>de</sub>	83.1 <sub>a</sub>	41	4 <sub>abcd</sub>	52.55
3	PHY 333 WRF	1264 <sub>ab</sub>	40.1 <sub>ab</sub>	4.6 <sub>gh</sub>	1.20 <sub>b</sub>	31.7 <sub>cde</sub>	82.3 <sub>bcd</sub>	41	4 <sub>bcd</sub>	54.75
4	NG 3405 B2XF	1264 <sub>ab</sub>	39.3 <sub>bcde</sub>	4.7 <sub>efg</sub>	1.11 <sub>i</sub>	28.9 <sub>g</sub>	81.2 <sub>e</sub>	31	3 <sub>g</sub>	55.95
5	PHY 444 WRF	1261 <sub>ab</sub>	39.8 <sub>bc</sub>	4.2 <sub>i</sub>	1.25 <sub>a</sub>	33.1 <sub>a</sub>	83.3 <sub>a</sub>	31	3 <sub>g</sub>	57.00
6	DG 3385 B2XF	1260 <sub>ab</sub>	39.7 <sub>bcd</sub>	5.1 <sub>b</sub>	1.15 <sub>gh</sub>	30.2 <sub>f</sub>	82.7 <sub>abc</sub>	31	3 <sub>g</sub>	54.10
7	ST 4946 GLB2	1257 <sub>abc</sub>	37.7 <sub>ghi</sub>	5.1 <sub>ab</sub>	1.17 <sub>def</sub>	33.0 <sub>ab</sub>	82.9 <sub>ab</sub>	31	4 <sub>def</sub>	52.90
8	DP 1518 B2XF	1246 <sub>abcd</sub>	38.7 <sub>def</sub>	4.6 <sub>h</sub>	1.17 <sub>de</sub>	30.5 <sub>f</sub>	82.4 <sub>bcd</sub>	41	4 <sub>bcd</sub>	54.50
9	DP 1612 B2XF	1244 <sub>abcd</sub>	37.8 <sub>fghi</sub>	4.9 <sub>cd</sub>	1.18 <sub>cd</sub>	32.3 <sub>bc</sub>	83.2 <sub>a</sub>	41	4 <sub>ab</sub>	54.85
10	ST 4848 GLT	1220 <sub>bcd</sub>	40.9 <sub>a</sub>	4.9 <sub>c</sub>	1.16 <sub>fgh</sub>	31.6 <sub>de</sub>	82.8 <sub>ab</sub>	31	4 <sub>def</sub>	55.15
11	NG 3406 B2XF	1208 <sub>bcde</sub>	38.9 <sub>cde</sub>	4.9 <sub>cd</sub>	1.14 <sub>h</sub>	30.1 <sub>f</sub>	82.6 <sub>abcd</sub>	31	3 <sub>fg</sub>	56.40
12	DP 1522 B2XF	1201 <sub>bcde</sub>	38.5 <sub>efg</sub>	5.1 <sub>ab</sub>	1.16 <sub>efg</sub>	31.2 <sub>e</sub>	82.6 <sub>abc</sub>	41	4 <sub>cde</sub>	52.35
13	ST 5032 GLT	1176 <sub>cde</sub>	37.2 <sub>i</sub>	4.6 <sub>h</sub>	1.21 <sub>b</sub>	32.8 <sub>ab</sub>	83.0 <sub>ab</sub>	31	4 <sub>cde</sub>	55.35
14	ST 4747 GLB2	1166 <sub>de</sub>	37.3 <sub>i</sub>	4.8 <sub>cde</sub>	1.20 <sub>b</sub>	31.4 <sub>de</sub>	82.0 <sub>cd</sub>	41	4 <sub>a</sub>	54.75
15	ST 5115 GLT	1129 <sub>e</sub>	37.5 <sub>hi</sub>	4.8 <sub>def</sub>	1.15 <sub>gh</sub>	31.9 <sub>cde</sub>	81.9 <sub>de</sub>	31	3 <sub>efg</sub>	56.50
<b>Mean</b>		<b>1233</b>	<b>38.8</b>	<b>4.8</b>	<b>1.18</b>	<b>31.5</b>	<b>82.6</b>	<b>31</b>	<b>4</b>	<b>54.79</b>
LSD ( $p \leq 0.05$ )		82	1.0	0.1	0.02	0.7	0.7		0.4	

\*Mean and LSD values were calculated from 15 varieties planted and harvested in fifteen independent 2016 Tennessee County Standard Trials.

## Replicated On-Farm Variety Trials

Lint yield, gin turnout, fiber quality and CCC loan value of 8 entries entered in the 2016 Cotton Incorporated Replicated Large Plot Variety Trial Program.\*\*

Yield Rank	Variety	Lint Yield (lb/ac)	Turnout (%)	Mic	Length (in.)	Strength (g/tex)	Unif. (%)	HVI Color	Leaf Grade	Loan Value (¢/lb)
1	PHY 444 WRF	1382 <sub>a</sub>	40.5 <sub>ab</sub>	4.3 <sub>e</sub>	1.25 <sub>a</sub>	33.0 <sub>a</sub>	83.4 <sub>a</sub>	31	3 <sub>cd</sub>	56.80
2	PHY 333 WRF	1342 <sub>ab</sub>	40.3 <sub>ab</sub>	4.8 <sub>d</sub>	1.20 <sub>b</sub>	32.2 <sub>b</sub>	82.5 <sub>b</sub>	31	4 <sub>a</sub>	56.70
3	ST 4946 GLB2	1326 <sub>ab</sub>	38.2 <sub>d</sub>	5.2 <sub>b</sub>	1.17 <sub>c</sub>	33.4 <sub>a</sub>	83.3 <sub>ab</sub>	31	3 <sub>bc</sub>	54.40
4	DG 3385 B2XF	1320 <sub>ab</sub>	39.8 <sub>bc</sub>	5.1 <sub>b</sub>	1.15 <sub>de</sub>	30.0 <sub>c</sub>	82.7 <sub>ab</sub>	31	3 <sub>d</sub>	54.60
5	DP 1614 B2XF	1314 <sub>ab</sub>	41.1 <sub>a</sub>	5.4 <sub>a</sub>	1.20 <sub>b</sub>	31.6 <sub>b</sub>	82.7 <sub>ab</sub>	31	4 <sub>a</sub>	52.95
6	ST 4848 GLT	1292 <sub>b</sub>	41.3 <sub>a</sub>	5.2 <sub>b</sub>	1.15 <sub>de</sub>	31.9 <sub>b</sub>	82.6 <sub>ab</sub>	31	4 <sub>a</sub>	54.25
7	NG 3406 B2XF	1270 <sub>b</sub>	38.7 <sub>cd</sub>	5.0 <sub>c</sub>	1.14 <sub>e</sub>	30.3 <sub>c</sub>	82.6 <sub>ab</sub>	31	3 <sub>bcd</sub>	56.25
8	DP 1522 B2XF	1266 <sub>b</sub>	38.9 <sub>cd</sub>	5.3 <sub>b</sub>	1.16 <sub>cd</sub>	31.5 <sub>b</sub>	82.8 <sub>ab</sub>	31	4 <sub>ab</sub>	54.25
<b>Mean</b>		<b>1313</b>	<b>39.8</b>	<b>5.0</b>	<b>1.18</b>	<b>31.7</b>	<b>82.8</b>	<b>31</b>	<b>3</b>	<b>55.03</b>
LSD ( $p \leq 0.05$ )		81	1.2	0.1	0.02	0.8	0.9		0.5	

\*\*Mean and LSD values were calculated from 8 entries replicated three times at four independent locations during the 2016 season.

# Official Variety Trial Results

## Commercial Varieties and Experimental Lines

Lint yield, gin turnout, and fiber quality of 32 entries entered in the 2016 Tennessee Official Variety Trial Program.\*

Yield Rank	Variety	Lint Yield (lb/ac)	Turnout (%)	Mic	Length (in.)	Strength (g/tex)	Unif. (%)	Leaf Grade
1	PHY 312 WRF	1533 a*	38.9 efg	4.7 klmn	1.17 cde	32.4 cde	82.8 abc	3 bcde
2	PHY 333 WRF	1529 a	40.4 abcd	4.7 j-n	1.16 defg	30.9 f-j	80.9 hijk	3 cdef
3	MON15R535 B2XF	1492 ab	41.3 a	4.9 e-k	1.15 e-j	30.6 g-k	81.2 g-k	2 ghi
4	CPS16214 B2XF	1466 abc	39.6 cde	4.5 no	1.23 a	31.0 f-j	81.5 e-i	3 defg
5	PHY 444 WRF	1465 abc	39.5 cde	4.4 o	1.24 a	33.3 bcd	82.8 abc	3 fghi
6	DP 1522 B2XF	1452 abcd	39.1 efg	5.4 a	1.13 jkl	30.4 hijk	82.1 b-g	3 defg
7	MON16R229 B2XF	1447 bcde	39.2 def	5.1 b-f	1.10 mn	30.5 hijk	80.9 hijk	3 defg
8	DP 1614 B2XF	1424 b-f	41.1 ab	5.3 ab	1.20 bc	31.4 efg	82.6 bcde	3 bcde
9	DG 3526 B2XF	1421 b-g	40.7 abc	4.9 f-l	1.14 ghij	29.6 k	82.4 b-f	3 defg
10	DG 3385 B2XF	1417 b-h	38.8 e-i	5.0 c-i	1.14 f-j	29.6 k	82.2 b-g	2 ghi
11	ST 4946 GLB2	1416 b-h	37.0 klm	4.9 d-i	1.13 ijkl	33.4 bc	81.8 c-i	3 cdef
12	DP 1518 B2XF	1405 c-i	37.7 hijk	4.6 mno	1.16 e-i	30.4 hijk	81.1 g-k	4 abc
13	BX 1771 GLTP	1399 c-i	37.8 hijk	5.0 b-h	1.14 f-j	33.7 ab	82.8 abc	3 cdef
14	BX 1737 GLT	1399 c-i	36.9 klm	4.7 i-n	1.17 def	31.3 e-i	82.0 b-g	3 fghi
15	AMX1601 B2XF	1397 c-i	40.0 bcde	5.1 b-g	1.18 bcd	34.1 ab	82.4 b-f	3 fghi
16	NG 3522 B2XF	1394 c-i	39.3 def	4.8 h-l	1.09 mn	27.9 l	80.8 ijk	2 hi
17	ST 4848 GLT	1379 d-j	39.0 efg	5.1 b-g	1.16 defg	31.5 efg	81.9 c-i	3 cdef
18	NG 3405 B2XF	1367 e-j	38.2 f-j	4.8 i-m	1.09 n	27.5 l	80.2 k	2 ghi
19	CG 3475 B2XF	1357 f-k	37.0 jklm	5.1 b-h	1.14 ghij	31.7 efg	82.9 abc	4 abcd
20	ST 4747 GLB2	1355 f-k	37.1 jklm	4.9 d-j	1.17 def	29.9 jk	80.3 k	3 cdef
21	SSG UA 222	1341 g-k	37.1 jklm	5.0 c-i	1.20 bc	32.3 cde	81.5 f-j	3 bcde
22	NG 3406 B2XF	1338 hijk	38.1 f-k	4.9 d-i	1.13 jkl	29.5 k	82.6 bcd	3 defg
23	DP 1612 B2XF	1330 ijkl	36.4 lm	5.0 b-h	1.16 defg	32.3 cde	83.0 ab	4 ab
24	BX 1775 GLTP	1330 ijkl	37.6 ijkl	4.7 klmn	1.17 defg	30.5 hijk	80.4 jk	2 hi
25	BX 1776 GLTP	1304 jkl	38.2 f-j	4.7 j-n	1.17 de	30.2 ijk	81.6 d-i	2 ghi
26	AMX1604 B2XF	1279 kl	37.9 g-k	5.1 bcde	1.13 hijk	32.4 cde	81.5 f-j	3 fghi
27	AMX1606 B2XF	1249 lm	36.3 mn	4.7 i-n	1.16 e-i	33.5 b	81.7 d-i	3 efg
28	SSG HQ 210	1194 mn	36.4 lm	5.2 abc	1.11 lmn	32.2 de	81.9 c-i	2 ghi
29	BRS 335	1168 mno	36.0 mn	4.6 lmno	1.16 efg	31.3 efg	81.2 ghijk	4 a
30	BRS 286	1145 no	35.1 no	4.9 g-l	1.11 klm	31.7 ef	81.9 c-h	3 defg
31	BRS 293	1116 no	34.6 o	5.2 abcd	1.13 ijkl	33.9 ab	82.3 b-f	2 i
32	DG 3544 B2XF	1098 o	37.1 jklm	5.1 b-g	1.20 b	34.8 a	83.8 a	3 fghi
<b>Average</b>		<b>1356</b>	<b>38.1</b>	<b>4.9</b>	<b>1.15</b>	<b>31.4</b>	<b>81.8</b>	<b>3</b>
LSD (p≤0.05)		82	1.3	0.3	0.03	1.1	1.1	0.7

\*Mean and LSD lint values were calculated from 32 entries replicated four times at five separate 2016 Tennessee Official Variety Trials. Mean and LSD fiber quality values were calculated from 32 representative samples from five 2016 Tennessee Official Variety Trials.

\*Means followed by the same letter are not significantly different.

## Two Year OVT Averages

Lint yield, gin turnout, and fiber quality of 19 like-entries entered in the 2015 and 2016 Tennessee Official Variety Trial Programs.\*

Yield Rank	Variety	Lint Yield (lb/ac)	Turnout (%)	Length		Strength		Leaf Grade
				Mic	(in.)	(g/tex)	Unif. (%)	
1	PHY 333 WRF	1525 <sub>a</sub>	39.9 <sub>ab</sub>	4.5 <sub>hi</sub>	1.18 <sub>def</sub>	31.1 <sub>efg</sub>	82.0 <sub>efg</sub>	4 <sub>cdef</sub>
2	PHY 312 WRF	1475 <sub>ab</sub>	38.5 <sub>cd</sub>	4.5 <sub>ghi</sub>	1.19 <sub>cd</sub>	32.1 <sub>cde</sub>	83.1 <sub>abcd</sub>	4 <sub>abcd</sub>
3	DP 1614 B2XF	1450 <sub>abc</sub>	40.7 <sub>a</sub>	5.0 <sub>ab</sub>	1.20 <sub>bc</sub>	31.1 <sub>efg</sub>	83.1 <sub>abc</sub>	4 <sub>b-f</sub>
4	PHY 444 WRF	1445 <sub>abc</sub>	39.0 <sub>bc</sub>	4.3 <sub>j</sub>	1.24 <sub>a</sub>	32.8 <sub>bc</sub>	83.2 <sub>abc</sub>	3 <sub>ghi</sub>
5	DP 1522 B2XF	1444 <sub>abc</sub>	38.7 <sub>cd</sub>	5.1 <sub>a</sub>	1.16 <sub>fghi</sub>	30.9 <sub>fgh</sub>	82.9 <sub>a-e</sub>	4 <sub>cdef</sub>
6	DG 3526 B2XF	1432 <sub>bcd</sub>	40.0 <sub>ab</sub>	4.8 <sub>cdef</sub>	1.16 <sub>fghi</sub>	30.1 <sub>h</sub>	82.9 <sub>abcd</sub>	4 <sub>efg</sub>
7	ST 4747 GLB2	1419 <sub>bcd</sub>	37.8 <sub>defg</sub>	4.7 <sub>cdefg</sub>	1.18 <sub>cde</sub>	30.4 <sub>gh</sub>	81.0 <sub>h</sub>	4 <sub>def</sub>
8	NG 3405 B2XF	1418 <sub>bcd</sub>	38.4 <sub>cd</sub>	4.6 <sub>fgh</sub>	1.12 <sub>j</sub>	28.0 <sub>i</sub>	81.3 <sub>gh</sub>	3 <sub>i</sub>
9	ST 4946 GLB2	1392 <sub>bcde</sub>	37.1 <sub>efgh</sub>	4.8 <sub>cdef</sub>	1.15 <sub>hi</sub>	33.4 <sub>ab</sub>	82.3 <sub>def</sub>	4 <sub>cdef</sub>
10	DP 1518 B2XF	1390 <sub>bcde</sub>	37.7 <sub>defg</sub>	4.4 <sub>ij</sub>	1.18 <sub>defg</sub>	30.4 <sub>gh</sub>	82.0 <sub>fg</sub>	5 <sub>abc</sub>
11	NG 3406 B2XF	1378 <sub>cde</sub>	38.0 <sub>cdef</sub>	4.7 <sub>c-g</sub>	1.15 <sub>hi</sub>	30.0 <sub>h</sub>	83.0 <sub>abcd</sub>	4 <sub>fgh</sub>
12	ST 4848 GLT	1373 <sub>cde</sub>	38.8 <sub>cd</sub>	4.8 <sub>b-f</sub>	1.17 <sub>d-h</sub>	31.3 <sub>efg</sub>	82.4 <sub>cdef</sub>	4 <sub>cdef</sub>
13	DP 1612 B2XF	1362 <sub>cde</sub>	36.9 <sub>fgh</sub>	4.8 <sub>bcde</sub>	1.18 <sub>cde</sub>	32.7 <sub>bcd</sub>	83.5 <sub>a</sub>	5 <sub>a</sub>
14	DG 3385 B2XF	1351 <sub>de</sub>	38.2 <sub>cde</sub>	4.8 <sub>b-f</sub>	1.17 <sub>d-h</sub>	30.5 <sub>gh</sub>	82.9 <sub>abcd</sub>	3 <sub>ghi</sub>
15	CG 3475 B2XF	1343 <sub>de</sub>	36.7 <sub>gh</sub>	4.9 <sub>bcd</sub>	1.16 <sub>e-i</sub>	31.7 <sub>def</sub>	83.3 <sub>ab</sub>	4 <sub>a-e</sub>
16	SSG UA 222	1302 <sub>ef</sub>	36.8 <sub>gh</sub>	4.7 <sub>d-h</sub>	1.21 <sub>b</sub>	31.9 <sub>cdef</sub>	82.0 <sub>efg</sub>	4 <sub>b-f</sub>
17	SSG HQ 210	1250 <sub>fg</sub>	36.1 <sub>hi</sub>	4.9 <sub>abc</sub>	1.14 <sub>ij</sub>	32.7 <sub>bcd</sub>	82.7 <sub>b-f</sub>	3 <sub>hi</sub>
18	BRS 335	1240 <sub>fg</sub>	36.1 <sub>hi</sub>	4.5 <sub>ghi</sub>	1.17 <sub>d-h</sub>	31.9 <sub>cdef</sub>	81.8 <sub>fgh</sub>	5 <sub>ab</sub>
19	BRS 293	1172 <sub>g</sub>	35.4 <sub>i</sub>	4.9 <sub>abcd</sub>	1.15 <sub>ghi</sub>	34.0 <sub>a</sub>	82.5 <sub>b-f</sub>	3 <sub>hi</sub>
20	BRS 286	1165 <sub>g</sub>	35.3 <sub>i</sub>	4.6 <sub>efgh</sub>	1.14 <sub>i</sub>	32.4 <sub>bcd</sub>	82.2 <sub>def</sub>	4 <sub>efg</sub>
<b>Average</b>		<b>1366</b>	<b>37.8</b>	<b>4.7</b>	<b>1.17</b>	<b>31.5</b>	<b>82.5</b>	<b>4</b>
LSD ( $p \leq 0.05$ )		90	1.1	0.2	0.02	1.1	0.8	0.6

\*Mean and LSD values were calculated from 20 identical-entries in the 2015 and 2016 Tennessee Official Variety Trials. Table compiled from UTIA AgResearch data of Raper et al. (2015) and Raper et al. (2016).

## Two Year CST Averages

Lint yield, gin turnout, and fiber quality of 10 like-entries in the 2015 and 2016 Tennessee County Standard Trial Programs.\*\*

Yield Rank	Variety	Lint Yield (lb/ac)	Turnout (%)	Mic	Length (in.)	Strength (g/tex)	Unif. (%)	HVI	Leaf Grade
								Color	
1	PHY 312 WRF	1325 <sub>a</sub>	38.1 <sub>b</sub>	4.5 <sub>cde</sub>	1.21 <sub>bc</sub>	32.4 <sub>b</sub>	83.5 <sub>bc</sub>	41	5 <sub>ab</sub>
2	PHY 444 WRF	1267 <sub>ab</sub>	39.5 <sub>a</sub>	4.0 <sub>f</sub>	1.27 <sub>a</sub>	33.1 <sub>a</sub>	84.0 <sub>a</sub>	31	4 <sub>e</sub>
3	PHY 333 WRF	1244 <sub>bc</sub>	39.6 <sub>a</sub>	4.5 <sub>cd</sub>	1.21 <sub>c</sub>	32.0 <sub>bc</sub>	83.0 <sub>cd</sub>	41	5 <sub>bc</sub>
4	DP 1518 B2XF	1242 <sub>bc</sub>	38.7 <sub>b</sub>	4.4 <sub>de</sub>	1.19 <sub>d</sub>	30.9 <sub>d</sub>	82.6 <sub>de</sub>	41	4 <sub>c</sub>
5	DG 3385 B2XF	1241 <sub>bc</sub>	39.4 <sub>a</sub>	4.9 <sub>a</sub>	1.16 <sub>f</sub>	30.5 <sub>d</sub>	83.3 <sub>bc</sub>	31	3 <sub>f</sub>
6	DP 1522 B2XF	1230 <sub>bcd</sub>	38.5 <sub>b</sub>	5.0 <sub>a</sub>	1.17 <sub>ef</sub>	31.6 <sub>c</sub>	83.3 <sub>bc</sub>	41	4 <sub>cd</sub>
7	ST 4946 GLB2	1221 <sub>bcd</sub>	37.5 <sub>c</sub>	4.9 <sub>a</sub>	1.18 <sub>de</sub>	33.2 <sub>a</sub>	83.5 <sub>ab</sub>	41	4 <sub>d</sub>
8	ST 4747 GLB2	1182 <sub>cd</sub>	37.2 <sub>c</sub>	4.6 <sub>b</sub>	1.21 <sub>bc</sub>	31.9 <sub>bc</sub>	82.5 <sub>de</sub>	41	5 <sub>a</sub>
9	ST 5032 GLT	1169 <sub>d</sub>	36.9 <sub>c</sub>	4.4 <sub>e</sub>	1.22 <sub>b</sub>	33.0 <sub>a</sub>	83.4 <sub>bc</sub>	41	4 <sub>c</sub>
10	ST 5115 GLT	1103 <sub>e</sub>	37.3 <sub>c</sub>	4.5 <sub>c</sub>	1.16 <sub>f</sub>	32.4 <sub>b</sub>	82.3 <sub>e</sub>	31	4 <sub>de</sub>
<b>Mean</b>		<b>1222</b>	<b>38.3</b>	<b>4.6</b>	<b>1.20</b>	<b>32.1</b>	<b>83.1</b>	<b>41</b>	<b>4</b>
LSD ( $p \leq 0.05$ )		64	0.6	0.1	0.01	0.6	0.5		0.4

\*\*Mean and LSD values were calculated from 10 identical-entries in the 2015 and 2016 Tennessee County Standard Trials. Table compiled from UTIA AgResearch data of Raper et al. (2015) and Raper et al. (2016).



For more information visit your county Extension Office or [utcrops.com](http://utcrops.com)



[AG.TENNESSEE.EDU](http://AG.TENNESSEE.EDU)